





Picture D1: Colours CMYNOLVA_m for homogeneity and goss images; PS operator $cmy0^* / 000n^* \text{ setcmykcolor}$



Picture D2: Radial gratings W-O, W-L, W-V, W-N, and W-Z; PS operator $cmy0^* \text{ setcmykcolor}$ (only)




Picture D3: 14 CIE-test colours and 2 + 16 grey steps; Use of PS operator $cmy0^* \text{ setcmykcolor}$ (only)


BAM-test chart no. BE16

Step: S2


Color reproduction properties of ISO/IEC TR 19797 and 24705




Picture D4: 16 equidistant steps W-O, W-L, W-V and W-N; Use of PS operator $cmy0^* \text{ setcmykcolor}$ (only)



Picture D5: Script and Landolt-rings N, O, L and V; Use of PS operator $cmy0^* \text{ setcmykcolor}$ (only)



Picture D6: Landolt-rings W-O and W-L; Use of PS operator $cmy0^* \text{ setcmykcolor}$ (only)



Picture D7: Landolt-rings W-V and W-N; Use of PS operator $cmy0^* \text{ setcmykcolor}$ (only)

input: $cmy0^* / 000n^* \text{ setcmykcolor}$

output: $cmy0^* / 000n^* \text{ setcmykcolor}$