






logarithmic N_a -sensitivity $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.38$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o + 0.39$
 Adaptation: $\lambda_{UT}=503$




logarithmic N_a -sensitivity $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.06$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o + 0.72$
 Adaptation: $\lambda_{UT}=525$




logarithmic N_a -sensitivity $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.79$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o - 0.02$
 Adaptation: $\lambda_{UT}=475$




logarithmic N_a -sensitivity $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 1.16$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o - 0.39$
 Adaptation: $\lambda_{UT}=450$




logarithmic N_a -saturation $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.38$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o + 0.39$
 Adaptation: $\lambda_{UT}=503$




logarithmic N_a -saturation $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.06$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o + 0.72$
 Adaptation: $\lambda_{UT}=525$




logarithmic N_a -saturation $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.79$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o - 0.02$
 Adaptation: $\lambda_{UT}=475$




logarithmic N_a -saturation $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 1.16$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o - 0.39$
 Adaptation: $\lambda_{UT}=450$




logarithmic N_a -sensitivity $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.38$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o + 0.39$
 Adaptation: $\lambda_{UT}=503$




logarithmic N_a -sensitivity $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.06$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o + 0.72$
 Adaptation: $\lambda_{UT}=525$




logarithmic N_a -sensitivity $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.79$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o - 0.02$
 Adaptation: $\lambda_{UT}=475$




logarithmic N_a -sensitivity $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 1.16$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o - 0.39$
 Adaptation: $\lambda_{UT}=450$




logarithmic N_a -saturation $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.38$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o + 0.39$
 Adaptation: $\lambda_{UT}=503$



logarithmic N_a -saturation $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.06$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o + 0.72$
 Adaptation: $\lambda_{UT}=525$



logarithmic N_a -saturation $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 0.79$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o - 0.02$
 Adaptation: $\lambda_{UT}=475$



logarithmic N_a -saturation $N_a = (\text{U}_a \cdot \text{T}_a)^{0.5}$ $\log N_a = \log N_o$
 $\log N_a = (\log \text{U}_a + \log \text{T}_a) / 2$ $\log \text{U}_a = \log \text{U}_o + 1.16$
 $\log [N_a, U_a, T_a]$ $\log \text{T}_a = \log \text{T}_o - 0.39$
 Adaptation: $\lambda_{UT}=450$

