

Optimal colours (o) *RYGCBM* of maximum (m) $C_{AB,10}$; D65, $Y_{me}=520_770$, LABHNU1_79

<i>Code, K=1:25</i>		L^*_{10}	A^*_{10}	B^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _o	570_770	64.15	64.78	55.73	85.45	0.1702	-0.0547	40.7	39	595	14	474
Y _{me}	520_770	86.53	7.14	78.71	79.03	0.1333	-0.0511	84.8	33	568	13	466
G _{me}	470_570	80.1	-68.74	39.71	79.39	0.0941	-0.0688	149.9	21	507	8	440
C _m	380_570	82.09	-43.23	-21.9	48.47	0.1078	-0.1006	206.8	15	477	44	621
B _{me}	380_520	56.52	-13.89	-63.53	65.03	0.1202	-0.1333	257.6	13	468	34	572
M _m	570_470	66.95	92.26	-33.68	98.22	0.1854	-0.1098	339.9	3	415	19	498
B _o	380_470	32.23	94.09	-92.25	131.77	0.2274	-0.1851	315.5	10	452	25	528
C _o	470_520	52.76	-61.75	0.52	61.76	0.0849	-0.0891	179.5	17	486	-1	486c
G _o	520_570	69.12	-47.05	59.68	75.99	0.1022	-0.0544	128.2	27	538	11	455
W	380_770	96.0	0.0	0.0	0.0	0.1298	-0.0895	0.0	-1	486c	17	486
N	380_770	22.33	0.0	0.0	0.0	0.1298	-0.0895	155.9	20	500	5	425