

Optimal colours (o) *RYGCBM* of maximum (m) $C_{AB,10}$; P60, $Y_m=510_770$, CIELAB_76

<i>Code</i>		L^*_{10}	a^*_{10}	b^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me}	570_770	66.33	67.03	114.37	132.57	0.2561	0.0	59.6	39	598	14	470
Y _{me}	510_770	93.01	-2.36	139.93	139.95	0.2143	-0.0219	90.9	33	566	13	465
G _{me}	470_570	82.02	-113.59	52.8	125.26	0.1575	-0.0592	155.0	22	513	-1	513c
C _m	380_570	84.09	-63.29	-27.42	68.97	0.1838	-0.0998	203.4	15	476	-1	476c
B _{me}	380_510	48.29	6.59	-88.23	88.48	0.2206	-0.1548	274.2	13	465	32	563
M _m	570_470	69.24	89.73	-39.0	97.84	0.2681	-0.109	336.5	-1	512c	22	512
R _o	570_440	67.0	80.66	4.26	80.77	0.264	-0.0836	3.0	-1	484c	16	484
G _o	520_570	74.34	-89.11	107.75	139.83	0.1661	-0.0265	129.5	27	536	9	448
W ₁	380_770	100.0	0.0	0.0	0.0	0.2154	-0.0861	0.0	-1	448c	-1	536c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) $C_{AB,10}$; P55, $Y_m=510_770$, CIELAB_76												
<i>Code</i>		L^*_{10}	a^*_{10}	b^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me}	570_770	67.2	67.44	115.87	134.06	0.256	0.0	59.7	39	597	14	470
Y _{me}	510_770	93.39	−0.82	139.56	139.56	0.2151	−0.0223	90.3	33	567	13	465
G _{me}	470_570	81.57	−112.65	50.71	123.54	0.1577	−0.0601	155.7	22	514	−1	514c
C _m	380_570	83.5	−66.3	−28.45	72.14	0.1821	−0.1004	203.2	15	476	−1	476c
B _{me}	380_510	47.17	2.43	−90.09	90.13	0.2174	−0.1575	271.5	13	465	33	566
M _m	570_470	69.83	87.65	−37.47	95.33	0.2665	−0.108	336.8	−1	513c	22	513
R _o	570_440	67.79	79.38	6.81	79.67	0.2628	−0.0821	4.9	−1	483c	16	483
G _o	520_570	74.16	−88.88	106.42	138.65	0.1661	−0.0271	129.8	27	537	9	447
W ₁	380_770	100.0	0.0	0.0	0.0	0.2154	−0.0861	0.0	−1	447c	−1	537c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) $C_{AB,10}$; P50, $Y_m=510_770$, CIELAB_76												
<i>Code</i>		L^*_{10}	a^*_{10}	b^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me}	570_770	68.23	67.66	117.65	135.72	0.2556	0.0	60.0	39	596	14	470
Y _{me}	510_770	93.82	0.77	139.04	139.05	0.2158	-0.0228	89.6	33	568	13	465
G _{me}	470_570	81.0	-111.65	48.15	121.59	0.1579	-0.0613	156.6	22	514	-1	514c
C _m	380_570	82.76	-69.78	-29.71	75.84	0.1801	-0.1012	203.0	15	476	-1	476c
B _{me}	380_510	45.83	-2.47	-92.33	92.36	0.2134	-0.1608	268.4	13	465	33	568
M _m	570_470	70.55	85.18	-35.57	92.31	0.2646	-0.1067	337.3	-1	514c	22	514
R _o	570_440	68.74	77.83	9.9	78.45	0.2614	-0.0803	7.2	-1	482c	16	482
G _o	520_570	73.92	-88.74	104.73	137.27	0.1661	-0.0279	130.2	27	538	9	445
W ₁	380_770	100.0	0.0	0.0	0.0	0.2154	-0.0861	0.0	-1	445c	-1	538c

Optimal colours (o) *RYGCBM* of maximum (m) $C_{AB,10}$; P45, $Y_m=510_770$, CIELAB_76

<i>Code</i>		L^*_{10}	a^*_{10}	b^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me}	570_770	69.48	67.58	119.79	137.54	0.255	0.0	60.5	39	595	14	470
Y _{me}	510_770	94.33	2.37	138.32	138.35	0.2165	-0.0234	89.0	33	569	13	465
G _{me}	470_570	80.25	-110.63	44.93	119.41	0.158	-0.0628	157.8	23	515	-1	515c
C _m	380_570	81.83	-73.79	-31.31	80.16	0.1777	-0.1021	202.9	15	477	-1	477c
B _{me}	380_510	44.19	-8.28	-95.05	95.41	0.2085	-0.1651	265.0	13	466	34	572
M _m	570_470	71.47	82.2	-33.17	88.64	0.2624	-0.1051	338.0	-1	515c	23	515
R _o	570_440	69.89	75.88	13.69	77.11	0.2596	-0.0782	10.2	-1	482c	16	482
G _o	520_570	73.56	-88.75	102.53	135.6	0.1659	-0.0289	130.8	27	539	8	443
W ₁	380_770	100.0	0.0	0.0	0.0	0.2154	-0.0861	0.0	-1	443c	-1	539c

Optimal colours (o) *RYGCBM* of maximum (m) $C_{AB,10}$; P40, $Y_m=510_770$, CIELAB_76

<i>Code</i>		L^*_{10}	a^*_{10}	b^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me}	570_770	71.0	66.97	122.42	139.54	0.2539	0.0	61.3	38	593	14	470
Y _{me}	510_770	94.91	3.89	137.29	137.35	0.2172	-0.0242	88.3	34	570	13	465
G _{me}	470_570	79.27	-109.6	40.79	116.95	0.1579	-0.0647	159.5	23	515	-1	515c
C _m	380_570	80.64	-78.37	-33.38	85.18	0.1749	-0.1034	203.0	15	477	-1	477c
B _{me}	380_510	42.15	-15.18	-98.42	99.58	0.2024	-0.1708	261.2	13	466	35	575
M _m	570_470	72.64	78.56	-30.08	84.12	0.2597	-0.1031	339.0	-1	516c	23	516
R _o	570_440	71.32	73.39	18.44	75.67	0.2574	-0.0756	14.1	-1	480c	16	480
G _o	520_570	73.04	-88.98	99.59	133.55	0.1655	-0.0302	131.7	28	540	7	439
W ₁	380_770	100.0	0.0	0.0	0.0	0.2154	-0.0861	0.0	-1	439c	-1	540c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) $C_{AB,10}$; P35, $Y_m=510_770$, CIELAB_76												
<i>Code</i>		L^*_{10}	a^*_{10}	b^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me}	570_770	72.89	65.5	125.67	141.72	0.2523	0.0	62.4	38	591	14	470
Y _{me}	510_770	95.58	5.17	135.76	135.86	0.2177	−0.0253	87.8	34	571	13	465
G _{me}	470_570	77.9	−108.58	35.33	114.18	0.1576	−0.0673	161.9	23	516	−1	516c
C _m	380_570	79.05	−83.51	−36.12	90.99	0.1715	−0.1051	203.3	15	478	−1	478c
B _{me}	380_510	39.57	−23.29	−102.66	105.27	0.1945	−0.1785	257.2	13	467	35	579
M _m	570_470	74.16	74.02	−25.98	78.45	0.2565	−0.1006	340.6	−1	517c	23	517
R _o	570_440	73.12	70.08	24.54	74.25	0.2547	−0.0724	19.3	−1	479c	15	479
G _o	520_570	72.24	−89.53	95.52	130.92	0.1647	−0.032	133.1	28	542	6	432
W ₁	380_770	100.0	0.0	0.0	0.0	0.2154	−0.0861	0.0	−1	432c	−1	542c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) $C_{AB,10}$; P30, $Y_m=510_770$, CIELAB_76												
<i>Code</i>		L^*_{10}	a^*_{10}	b^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me}	570_770	75.28	62.63	129.8	144.12	0.2497	0.0	64.2	38	590	13	469
Y _{me}	510_770	96.36	5.99	133.38	133.52	0.2181	-0.0268	87.4	34	573	13	465
G _{me}	470_570	75.95	-107.52	27.85	111.07	0.157	-0.071	165.4	23	516	-1	516c
C _m	380_570	76.84	-89.02	-39.92	97.56	0.1675	-0.1076	204.1	15	478	-1	478c
B _{me}	380_510	36.21	-32.63	-108.14	112.96	0.1841	-0.1897	253.2	13	468	36	583
M _m	570_470	76.19	68.2	-20.4	71.19	0.2524	-0.0972	343.3	-1	519c	23	519
R _o	570_440	75.44	65.52	32.58	73.17	0.2513	-0.0683	26.4	-1	478c	15	478
G _o	520_570	70.98	-90.47	89.63	127.35	0.1634	-0.0346	135.2	28	544	2	414
W ₁	380_770	100.0	0.0	0.0	0.0	0.2154	-0.0861	0.0	-1	414c	-1	544c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) $C_{AB,10}$; P25, $Y_m=510_770$, CIELAB_76												
<i>Code</i>		L^*_{10}	a^*_{10}	b^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me}	570_770	78.39	57.48	135.15	146.87	0.2459	0.0	66.9	37	588	13	469
Y _{me}	510_770	97.25	6.04	129.48	129.62	0.2181	-0.029	87.3	35	576	12	464
G _{me}	470_570	73.01	-106.24	17.09	107.6	0.1557	-0.0765	170.8	23	515	-1	515c
C _m	380_570	73.63	-94.33	-45.45	104.71	0.1628	-0.1115	205.7	15	479	-1	479c
B _{me}	380_510	31.68	-42.8	-115.39	123.07	0.1705	-0.2072	249.6	13	468	37	586
M _m	570_470	78.94	60.51	-12.44	61.78	0.2473	-0.0927	348.3	-1	518c	23	518
R _o	570_440	78.47	58.99	43.6	73.35	0.2466	-0.0631	36.4	58	693	15	475
G _o	520_570	68.89	-91.77	80.59	122.14	0.1614	-0.0387	138.7	29	547	-1	547c
W ₁	380_770	100.0	0.0	0.0	0.0	0.2154	-0.0861	0.0	-1	475c	-1	547c