

$XYZ_W = 103.66, 99.99, 52.43$

$A_2 = 2.5 (a_2 - a_{2,n}) Y$

$B_2 = 2.5 B_c (b_2 - b_{2,n}) Y$

$a_2 = a_{20} [(x - x_c) / y]$

$b_2 = b_{20} [z / y]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 1.800$

$C_{AB2} = [A_2^2 + B_2^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) C_{AB} in

linear colour space ($C_{AB,2n}, Y$)

Illumin. P35, $Y_W = 100, Y_N = 4$

Name	Range	X_d	Y_d	Z_d	x_d	y_d	λ_d	λ_c
R_d	575_775	71.28	40.36	2.18	0.6262	0.3545	605	496
Y_d	500_775	95.76	94.74	4.5	0.491	0.4858	578	472
G_d	500_575	28.72	58.47	4.46	0.3133	0.6379	548	548c
C_d	380_575	36.63	63.73	52.4	0.2397	0.4172	496	605
B_d	380_500	12.15	9.35	50.08	0.1698	0.1307	472	578
M_d	575_500	79.19	45.62	50.11	0.4527	0.2607	548c	548
W_d	380_775	103.66	99.99	52.43	0.4047	0.3904	100%	
N_d	380_775	4.14	3.99	2.09	0.4047	0.3904	4%	
Z_d	380_775	18.66	18.0	9.43	0.4047	0.3904	18%	

$L^* = 60 \log[f(Y_{an})]$



$f(Y_{an}) = \pm [1 + 10 |Y_{an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y & Name

Illuminant P35

$Y_W = 100, Y_N = 4$