

$XYZ_W = 97.45, 100.0, 95.98$

$A_2 = 2.5 (a_2 - a_{2,n}) Y$

$B_2 = 2.5 B_c (b_2 - b_{2,n}) Y$

$a_2 = a_{20} [(x - x_c) / y]$

$b_2 = b_{20} [z / y]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 0.900$

$C_{AB2} = [A_2^2 + B_2^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) C_{AB} in

linear colour space ($C_{AB,2}, Y$)

Illumin. P55, $Y_W = 100, Y_N = 4$

Name	Range	X_d	Y_d	Z_d	x_d	y_d	λ_d	λ_c
R_d	569_775	65.4	42.64	3.98	0.5837	0.3806	597	490
Y_d	494_775	82.28	94.94	9.62	0.4403	0.5081	572	464
G_d	494_569	20.87	56.4	9.57	0.2403	0.6493	536	536c
C_d	380_569	36.05	61.46	95.94	0.1863	0.3176	490	597
B_d	380_494	19.17	9.15	90.3	0.1616	0.0771	464	572
M_d	569_494	80.57	47.69	90.35	0.3685	0.2181	536c	536
W_d	380_775	97.45	100.0	95.98	0.3321	0.3407	100%	
N_d	380_775	3.89	4.0	3.83	0.3321	0.3407	4%	
Z_d	380_775	17.54	18.0	17.27	0.3321	0.3407	18%	

$L^* = 60 \log[f(Y_{an})]$



$f(Y_{an}) = \pm [1 + 10 |Y_{an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y & Name

Illuminant P55

$Y_W = 100, Y_N = 4$