

$XYZ_W = 97.45, 100.0, 95.98$

$A_2 = 2.5 (a_2 - a_{2,n}) Y$

$B_2 = 2.5 B_c (b_2 - b_{2,n}) Y$

$a_2 = a_{20} [(x - x_c) / y]$

$b_2 = b_{20} [z / y]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 0.900$

$C_{AB2} = [A_2^2 + B_2^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) C_{AB} in

linear colour space ($C_{AB,2}, Y$)

Illumin. P55, $Y_W = 100, Y_N = 10$

Name	Range	X_d	Y_d	Z_d	x_d	y_d	λ_d	λ_c
R_d	569_775	67.41	46.23	9.73	0.5463	0.3747	597	490
Y_d	494_775	83.23	95.26	15.02	0.43	0.4922	572	464
G_d	494_569	25.67	59.13	14.98	0.2572	0.5926	536	536c
C_d	380_569	39.9	63.87	95.94	0.1997	0.3198	490	597
B_d	380_494	24.07	14.84	90.66	0.1857	0.1145	464	572
M_d	569_494	81.63	50.97	90.7	0.3655	0.2282	536c	536
W_d	380_775	97.45	100.0	95.98	0.3321	0.3407	100%	
N_d	380_775	9.74	10.0	9.59	0.3321	0.3407	10%	
Z_d	380_775	17.54	18.0	17.27	0.3321	0.3407	18%	

$L^* = 60 \log[f(Y_{an})]$



$f(Y_{an}) = \pm [1 + 10 |Y_{an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y & Name

Illuminant P55

$Y_W = 100, Y_N = 10$