

$XYZ_W = 97.93, 100.0, 118.95$

$A_2 = 2.5 (a_2 - a_{2,n}) Y$

$B_2 = 2.5 B_c (b_2 - b_{2,n}) Y$

$a_2 = a_{20} [(x - x_c) / y]$

$b_2 = b_{20} [z / y]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 0.700$

$C_{AB2} = [A_2^2 + B_2^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) C_{AB} in

linear colour space ($C_{AB,2}, Y$)

Illumin. Q00, $Y_W = 100, Y_N = 0$

Name	Range	X_d	Y_d	Z_d	x_d	y_d	λ_d	λ_c
R_d	567_775	61.05	39.04	0.28	0.6081	0.3889	596	487
Y_d	492_775	77.98	94.36	7.21	0.4343	0.5255	570	462
G_d	492_567	17.13	55.52	7.16	0.2146	0.6956	535	535c
C_d	380_567	37.07	61.15	118.9	0.1707	0.2816	487	596
B_d	380_492	20.14	5.83	111.98	0.1459	0.0422	462	570
M_d	567_492	80.99	44.67	112.03	0.3407	0.1879	535c	535
W_d	380_775	97.93	100.0	118.95	0.309	0.3155	100%	
N_d	380_775	0.09	0.1	0.11	0.3089	0.3154	0%	
Z_d	380_775	17.62	18.0	21.41	0.309	0.3155	18%	

$L^* = 60 \log[f(Y_{an})]$



$f(Y_{an}) = \pm [1 + 10 |Y_{an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y & Name

Illuminant Q00

$Y_W = 100, Y_N = 0$