

$$Y_{adj} = [Y - 50] / 50$$

$$A_2 = 2,5 (a_2 - a_{2,n}) Y$$

$$B_2 = 2,5 B_c (b_2 - b_{2, \text{н}}) Y$$

$$a_2 = a_{20} [(x - x_c) / y]$$

$$b_2 = b_{20} [x/y]$$

$$a_{20} = 1, b_{20} = -0,4$$

$$x_c = 0,110, B_c = 0,800$$

$$C_{AB2} = [A_2^2 + B_2^2]^{1/2}$$

6 Ostwald colours (o)

of maximum (m) C_{A0} in

linear colour space ($C_{AB,2}, Y$)

Illumin. D65, $Y_{ref}=100$, $Y_N=10$

Name	Range	X_d	Y_d	Z_d	x_d	y_d	λ_d	λ_c
R_d	567_775	63.21	44.18	11.04	0.5337	0.373	596	489
Y_d	493_775	78.93	94.84	17.06	0.4135	0.4969	570	463
G_d	493_567	25.32	60.76	17.02	0.2455	0.5892	535	535c
C_d	380_567	41.44	65.92	108.85	0.1916	0.3048	489	596
B_d	380_493	25.71	15.26	102.83	0.1788	0.1061	463	570
M_d	567_493	79.33	49.34	102.87	0.3426	0.2131	535c	535
W_d	380_775	95.04	100.0	108.89	0.3127	0.329		100%
N_d	380_775	9.5	10.0	10.88	0.3127	0.329		10%
Z_d	380_775	17.1	18.0	19.6	0.3127	0.329		18%

$$f(Y_{20}) = \pm [1 + 10 |Y_{20}|^n]$$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y & Name

Illuminant D65

$$Y_W=100, Y_N=10$$
