

$XYZ_{W,10} = 116.54, 99.99, 26.13$

$A_{2,10} = 2.5 (a_{2,10} - a_{2,n,10}) Y_{10}$

$B_{2,10} = 2.5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$

$a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$

$b_{2,10} = b_{20} [z_{10} / y_{10}]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 3.700$

$C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) $C_{AB,10}$ in

linear colour space ($C_{AB,2,10}, Y_{10}$)

Illumin. P25, $Y_{W,10} = 100, Y_{N,10} = 4$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	λ_d	λ_c
R_d	578_775	87.32	48.86	1.07	0.6362	0.3559	605	495
Y_d	500_775	112.7894	86	2.25	0.5373	0.4519	580	473
G_d	500_578	30.23	50.1	2.25	0.366	0.6066	547	547c
C_d	380_578	33.99	55.24	26.13	0.2946	0.4787	495	605
B_d	380_500	8.54	9.23	24.95	0.1999	0.2161	473	580
M_d	578_500	91.09	54.0	24.95	0.5356	0.3175	547c	547
W_d	380_775	116.5499	99.99	26.13	0.4802	0.412	100%	
N_d	380_775	4.66	3.99	1.04	0.4802	0.412	4%	
Z_d	380_775	20.97	17.99	4.7	0.4802	0.412	18%	

$L^*_{10} = 60 \log[f(Y_{10,an})]$ $Y_{10,an} = [Y_{10} - 50] / 50$



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:
 Y_{10} & Name
 Illuminant P25
 $Y_{W,10} = 100, Y_{N,10} = 4$