

$XYZ_{W,10}=109.29, 99.99, 39.19$

$A_{2,10} = 2,5 (a_{2,10} - a_{2,n,10}) Y_{10}$

$B_{2,10} = 2,5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$

$a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$

$b_{2,10} = b_{20} [z_{10} / y_{10}]$

$a_{20} = 1, b_{20} = -0,4$

$x_c = 0,110, B_c = 2,500$

$C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$


6 Ostwald colours (o)

of maximum (m) $C_{AB,10}$ in

linear colour space ($C_{AB,2,10}, Y_{10}$)

Illumin. P30, $Y_{W,10}=100, Y_{N,10}=10$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	λ_d	λ_c
R_d	574_775	82.51	50.69	3.95	0.6015	0.3695	602	491
Y_d	497_775	103.81	95.02	5.9	0.507	0.4641	577	469
G_d	497_574	32.33	54.43	5.9	0.3488	0.5873	542	542c
C_d	380_574	37.82	59.41	39.2	0.2772	0.4354	491	602
B_d	380_497	16.53	15.08	37.25	0.24	0.219	469	577
M_d	574_497	88.01	55.67	37.25	0.4864	0.3077	542c	542
W_d	380_775	109.29	99.99	39.19	0.4398	0.4024	100%	
N_d	380_775	10.92	9.99	3.91	0.4398	0.4024	10%	
Z_d	380_775	19.67	17.99	7.05	0.4398	0.4024	18%	



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$
 n increases to 1 for:
 1. decreasing of the contrast C
 2. adjacent compared to separate colours.

Parameter:
 Y_{10} & Name
 Illuminant P30
 $Y_{W,10}=100, Y_{N,10}=10$