

$XYZ_{W,10} = 104.71, 99.99, 52.16$

$A_{2,10} = 2.5 (a_{2,10} - a_{2,n,10}) Y_{10}$

$B_{2,10} = 2.5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$

$a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$

$b_{2,10} = b_{20} [z_{10} / y_{10}]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 1.800$

$C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) $C_{AB,10}$ in

linear colour space ($C_{AB,2,10}, Y_{10}$)

Illumin. P35, $Y_{W,10} = 100, Y_{N,10} = 50$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	λ_d	λ_c
R_d	571_775	88.62	70.8	26.13	0.4775	0.3815	601	489
Y_d	494_775	100.6	197.16	27.78	0.446	0.4307	574	466
G_d	494_571	64.44	76.45	27.78	0.382	0.4532	540	540c
C_d	380_571	68.6	79.34	52.18	0.3427	0.3964	489	601
B_d	380_494	56.61	52.98	50.53	0.3535	0.3308	466	574
M_d	571_494	92.78	73.69	50.53	0.4275	0.3395	540c	540
W_d	380_775	104.7	199.99	52.16	0.4076	0.3892	100%	
N_d	380_775	52.35	49.99	26.08	0.4076	0.3892	50%	
Z_d	380_775	18.84	17.99	9.38	0.4076	0.3892	18%	

$L^*_{10} = 60 \log[f(Y_{10,an})]$ $Y_{10,an} = [Y_{10} - 50] / 50$



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y_{10} & Name

Illuminant P35

$Y_{W,10} = 100, Y_{N,10} = 50$