

$XYZ_{W,10} = 109.29, 99.99, 39.19$

$A_{2,10} = 2,5 (a_{2,10} - a_{2,n,10}) Y_{10}$

$B_{2,10} = 2,5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$

$a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$

$b_{2,10} = b_{20} [z_{10} / y_{10}]$

$a_{20} = 1, b_{20} = -0,4$

$x_c = 0,110, B_c = 2,500$

$C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$

6 Ostwald colours (o)

of maximum (m) $C_{AB,10}$ in


linear colour space ($C_{AB,2,10}, Y_{10}$)

Illumin. P30, $Y_{W,10} = 100, Y_{N,10} = 50$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	λ_d	λ_c
R_d	574_775	94.46	72.65	19.63	0.5058	0.389	602	491
Y_d	497_775	106.2999	72.8	20.72	0.4739	0.4337	577	469
G_d	497_574	66.58	74.73	20.72	0.4109	0.4611	542	542c
C_d	380_574	69.63	77.49	39.21	0.3736	0.4158	491	602
B_d	380_497	57.8	52.86	38.13	0.3884	0.3552	469	577
M_d	574_497	97.51	75.41	38.13	0.462	0.3573	542c	542
W_d	380_775	109.2999	99.99	39.19	0.4398	0.4024	100%	
N_d	380_775	54.64	49.99	19.59	0.4398	0.4024	50%	
Z_d	380_775	19.67	17.99	7.05	0.4398	0.4024	18%	

$L^*_{10} = 60 \log[f(Y_{10,an})]$

$Y_{10,an} = [Y_{10} - 50] / 50$



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$
 n increases to 1 for:
 1. decreasing of the contrast C
 2. adjacent compared to separate colours.

Parameter:
 Y_{10} & Name
 Illuminant P30
 $Y_{W,10} = 100, Y_{N,10} = 50$