

Transformation between the *Judd* tristimulus and opponent values

Data see K. Richter, PhD thesis, University of Basel (Switzerland), 1969, page 58.

elementary colour	dominant wavelength	Judd spectral tristimulus values			chromatic values	
		$\bar{x}(\lambda)$	$\bar{y}(\lambda)$	$\bar{z}(\lambda)$	$\bar{a}(\lambda)$	$\bar{b}(\lambda)$
blue	$\lambda_B=475$ nm	0,8267	0,9339	0,0017	0,0000	–
green	$\lambda_G=502$ nm	0,0107	0,0038	0,0005	–1,0000	0,0000
yellow	$\lambda_Y=574$ nm	0,1304	0,1124	0,9281	0,0000	1,0000
red	$\lambda_R=494$ c,E nm	0,0028	0,3701	0,2238	–	0,0000

There are six equations to calculate the six constants: b_{21} to b_{33}

$$\bar{a}(\lambda_B)=b_{21}\bar{x}(\lambda_B)+b_{22}\bar{y}(\lambda_B)+b_{23}\bar{z}(\lambda_B)=0 \quad \bar{b}(\lambda_G)=b_{31}\bar{x}(\lambda_G)+b_{32}\bar{y}(\lambda_G)+b_{33}\bar{z}(\lambda_G)=0$$

$$\bar{a}(\lambda_G)=b_{21}\bar{x}(\lambda_G)+b_{22}\bar{y}(\lambda_G)+b_{23}\bar{z}(\lambda_G)=-1 \quad \bar{b}(\lambda_Y)=b_{31}\bar{x}(\lambda_Y)+b_{32}\bar{y}(\lambda_Y)+b_{33}\bar{z}(\lambda_Y)=1$$

$$\bar{a}(\lambda_Y)=b_{21}\bar{x}(\lambda_Y)+b_{22}\bar{y}(\lambda_Y)+b_{23}\bar{z}(\lambda_Y)=0 \quad \bar{b}(\lambda_R)=b_{31}\bar{x}(\lambda_R)+b_{32}\bar{y}(\lambda_R)+b_{33}\bar{z}(\lambda_R)=0$$

Together with the use of the standard equation: $\bar{l}(\lambda)=\bar{y}(\lambda)$ (1)

the equations between spectral opponent and tristimulus colour values are:

$$\begin{pmatrix} \bar{l}(\lambda) \\ \bar{a}(\lambda) \\ \bar{b}(\lambda) \end{pmatrix} = \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{pmatrix} \cdot \begin{pmatrix} \bar{x}(\lambda) \\ \bar{y}(\lambda) \\ \bar{z}(\lambda) \end{pmatrix} = \begin{pmatrix} 0,0000 & 1,0000 & 0,0000 \\ 2,9797 & -2,6662 & -0,0960 \\ -0,4139 & 1,4571 & -2,4046 \end{pmatrix} \cdot \begin{pmatrix} \bar{x}(\lambda) \\ \bar{y}(\lambda) \\ \bar{z}(\lambda) \end{pmatrix} \quad (2)$$

Remark: The weighting ratio in the RG and YB direction is 2,8:1 or 1:0,3571.