

$XYZ_w = 84.1998, 88.59, 96.46$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2191$

$b_2 = -[1/Z_n]^{1/3} = -0.08376$

$n = D65$

CIELAB 76

Nome e gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald i colori ottimali, $Y_w = 88,6$

di massima (m) C_{AB} sotto illuminante D65

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 84.1998, 88.59, 96.46$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2191$

$b_2 = -[1/Z_n]^{1/3} = -0.08376$

$n = D65$

CIELAB 76

Nome e gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald i colori ottimali, $Y_w = 88,6$

di massima (m) C_{AB} sotto illuminante D65

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 84.1998, 88.59, 96.46$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2191$

$b_2 = -[1/Z_n]^{1/3} = -0.08376$

$n = D65$

CIELAB 76

Nome e gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald i colori ottimali, $Y_w = 88,6$

di massima (m) C_{AB} sotto illuminante D65

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 84.1998, 88.59, 96.46$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2191$

$b_2 = -[1/Z_n]^{1/3} = -0.08376$

$n = D65$

CIELAB 76

Nome e gamma spettrale

R_m 561_770 Y_m 520_770

G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald i colori ottimali, $Y_w = 88,6$

di massima (m) C_{AB} sotto illuminante D65

nel diagramma di CIELAB (a^*, b^*)

