

$$H^*_e = \text{G50B}_e$$

Data for maksimalfarge (Ma):

LabCh*_{e, Ma}: 56 -39 -29 49 216

***HIC**_{e, Ma}: G50B_100_100_e**

*rgbic**_{e, Ma}:

0.0 1.0 0.73 1.0 1.0

trekantslyshet T^*

%Omfang

$$u^*_{rel} = 92$$


%Regularitet

$$g^*_{H,rel} = 57$$
 $\delta^{13}\text{C}_{\text{rel}} = 50$

Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains.

I:

THE



ORS20a; adapterte (a) CIELAB data					
navn	L^*_a	a^*_a	b^*_a	$C^*_{ab,a}$	$h^*_{ab,a}$
R _{e, Ma}	47.6	64.9	30.9	71.9	25
Y _{e, Ma}	82.9	-3.5	87.8	87.9	92
G _{e, Ma}	52.4	-67.1	21.5	70.5	162
C _{e, Ma}	56.6	-39.7	-29.9	49.8	216
B _{e, Ma}	37.9	1.3	-45.4	45.4	271
M _{e, Ma}	34.8	49.2	-30.0	57.7	328
N _{e, Ma}	17.7	0.0	0.0	0.0	0
W _{e, Ma}	95.4	0.0	0.0	0.0	0
R _{e, CIE}	39.9	58.7	27.9	65.0	25
Y _{e, CIE}	81.2	-2.8	71.5	71.6	92
G _{e, CIE}	52.2	-42.4	13.6	44.5	162
B _{e, CIE}	30.5	1.4	-46.4	46.4	271

6 ORS20a; adapterte (a) CIELAB data					
H_e^*	$L^*=L_a^*a^*$	b_a^*	$C_{ab,a}^*$	$h_{ab,a}^*$	
R00Y_100_100 _e	47.6	64.9	30.9	71.9	25
R25Y_100_100 _e	51.5	54.2	47.2	71.9	41
R50Y_100_100 _e	60.3	35.6	59.0	68.9	58
R75Y_100_100 _e	70.4	17.0	72.2	74.1	76
Y00G_100_100 _e	82.9	-3.5	87.8	87.9	92
Y25G_100_100 _e	76.9	-25.5	75.9	80.1	108
Y50G_100_100 _e	65.8	-41.4	54.4	68.3	127
Y75G_100_100 _e	56.9	-56.3	38.1	68.0	145
G00B_100_100 _e	52.4	-67.1	21.5	70.5	162
G25B_100_100 _e	54.6	-53.2	-9.0	53.9	189
G50B_100_100 _e	56.6	-39.7	-29.9	49.8	216
G75B_100_100 _e	52.7	-21.1	-44.1	48.9	244
B00R_100_100 _e	37.9	1.3	-45.4	45.4	271
B25R_100_100 _e	26.7	26.6	-45.8	52.9	300
B50R_100_100 _e	34.8	49.2	-30.0	57.7	328
B75R_100_100 _e	47.3	71.5	-9.9	72.1	352

