

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for D65, $Y_w=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	70.18	61.64	132.51	146.14	0.2511	−0.0093	65.0	37 589	15 475
Y _m 495_770	97.55	−23.15	119.09	121.32	0.2052	−0.0337	101.0	32 563	14 470
G _o 495_565	77.76	−132.05	85.19	157.15	0.145	−0.0407	147.1	25 525	−1 525c
C _m 380_565	81.29	−66.98	−32.23	74.33	0.181	−0.1027	205.6	16 481	−1 481c
B _m 380_495	29.91	89.06	−117.04	147.07	0.3122	−0.2135	307.2	12 460	28 543
M _o 575_475	74.32	82.25	−40.48	91.68	0.2609	−0.1085	333.7	−1 519c	23 519
N _o 380_770	0.9	−0.01	0.0	0.01	0.2154	−0.0861	158.2	23 515	−1 515c
W _o 380_770	100.0	0.0	0.0	0.0	0.2154	−0.0861	0.0	−1 519c	−1 515c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for D50, $Y_w=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	72.41	64.43	122.61	138.51	0.2528	−0.0092	62.2	38 590	15 476
Y _m 495_770	98.02	−15.61	105.1	106.25	0.2096	−0.0324	98.4	33 566	14 470
G _o 495_565	76.49	−129.88	71.52	148.28	0.1462	−0.0399	151.1	25 525	−1 525c
C _m 380_565	79.46	−79.35	−32.24	85.65	0.1749	−0.0954	202.1	16 483	−1 483c
B _m 380_495	26.85	73.87	−110.83	133.19	0.3025	−0.2077	303.6	12 462	29 549
M _o 575_475	75.64	77.8	−34.15	84.96	0.2588	−0.0971	336.2	−1 520c	24 520
N _o 380_770	0.9	−0.01	0.0	0.01	0.2164	−0.0785	160.0	23 516	−1 516c
W _o 380_770	100.0	0.0	0.0	0.0	0.2164	−0.0785	0.0	−1 520c	−1 516c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for P40, $Y_w=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	75.32	62.78	115.6	131.55	0.2541	−0.0091	61.4	38 591	15 476
Y _m 495_770	98.43	−11.1	94.89	95.53	0.2149	−0.0309	96.6	33 569	14 470
G _o 495_565	74.32	−130.17	60.25	143.44	0.1477	−0.039	155.1	25 525	−1 525c
C _m 380_565	76.81	−88.65	−33.56	94.79	0.172	−0.0905	200.7	16 484	−1 484c
B _m 380_495	23.71	63.97	−106.28	124.05	0.3002	−0.2061	301.0	12 462	30 554
M _o 575_475	77.75	72.17	−27.97	77.4	0.2582	−0.0873	338.8	−1 521c	24 521
N _o 380_770	0.9	−0.01	0.0	0.01	0.2197	−0.0724	161.7	23 518	−1 518c
W _o 380_770	100.0	0.0	0.0	0.0	0.2197	−0.0724	0.0	−1 521c	−1 518c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for A00, $Y_w=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	79.72	59.98	96.44	113.57	0.2573	−0.0089	58.1	38 592	15 477
Y _m 495_770	99.03	−4.25	71.77	71.89	0.2242	−0.0281	93.3	35 575	14 470
G _o 495_565	70.4	−130.5	38.18	135.97	0.1505	−0.0372	163.6	25 525	−1 525c
C _m 380_565	72.11	−107.42	−33.03	112.38	0.1651	−0.078	197.0	17 488	−1 488c
B _m 380_495	17.93	38.27	−92.55	100.15	0.2824	−0.1957	292.4	12 464	33 566
M _o 575_475	81.12	63.13	−17.56	65.53	0.2585	−0.0683	344.4	−1 524c	24 524
N _o 380_770	0.9	−0.01	0.0	0.01	0.226	−0.0593	165.2	24 523	−1 523c
W _o 380_770	100.0	0.0	0.0	0.0	0.226	−0.0593	0.0	−1 524c	−1 523c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for E00, $Y_w=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	72.66	59.89	132.18	145.12	0.2528	−0.0092	65.6	38 590	15 475
Y _m 495_770	97.85	−20.53	116.26	118.06	0.21	−0.0327	100.0	33 566	14 470
G _o 495_565	76.01	−134.83	79.92	156.74	0.1458	−0.0403	149.3	25 525	−1 525c
C _m 380_565	79.25	−71.99	−34.73	79.94	0.1813	−0.1019	205.7	16 481	−1 481c
B _m 380_495	28.02	86.89	−116.93	145.68	0.3176	−0.2164	306.6	12 460	29 546
M _o 575_475	76.13	78.35	−36.32	86.36	0.2615	−0.1034	335.1	−1 520c	24 520
N _o 380_770	0.9	−0.01	0.0	0.01	0.2191	−0.0837	159.0	23 516	−1 516c
W _o 380_770	100.0	0.0	0.0	0.0	0.2191	−0.0837	0.0	−1 520c	−1 516c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for C00, $Y_w=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	70.87	57.84	137.7	149.35	0.2509	−0.0093	67.2	37 589	15 475
Y _m 495_770	97.36	−24.85	124.71	127.16	0.2067	−0.0335	101.2	32 564	14 470
G _o 495_565	76.89	−132.62	88.67	159.53	0.1463	−0.0408	146.2	25 526	−1 526c
C _m 380_565	80.74	−63.1	−34.08	71.72	0.185	−0.1061	208.3	16 480	−1 480c
B _m 380_495	31.05	90.91	−118.66	149.48	0.3141	−0.2145	307.4	12 460	28 543
M _o 575_475	75.24	80.09	−40.36	89.69	0.2615	−0.1106	333.2	−1 519c	23 519
N _o 380_770	0.9	−0.01	0.0	0.01	0.2176	−0.0885	157.8	23 515	−1 515c
W _o 380_770	100.0	0.0	0.0	0.0	0.2176	−0.0885	0.0	−1 519c	−1 515c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for P00, $Y_w=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	74.58	60.91	124.91	138.97	0.2542	−0.0091	64.0	38 591	15 475
Y _m 495_770	98.18	−15.24	106.16	107.25	0.2139	−0.0316	98.1	33 568	14 470
G _o 495_565	74.68	−133.58	69.73	150.69	0.1469	−0.0396	152.4	25 525	−1 525c
C _m 380_565	77.52	−81.23	−35.08	88.48	0.1771	−0.0968	203.3	16 483	−1 483c
B _m 380_495	25.68	75.68	−112.26	135.38	0.3112	−0.2126	303.9	12 461	30 551
M _o 575_475	77.42	74.31	−31.41	80.68	0.2603	−0.0948	337.0	−1 521c	24 521
N _o 380_770	0.9	−0.01	0.0	0.01	0.2206	−0.078	160.5	23 517	−1 517c
W _o 380_770	100.0	0.0	0.0	0.0	0.2206	−0.078	0.0	−1 521c	−1 517c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for Q00, $Y_w=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	70.65	58.47	137.7	149.6	0.2513	−0.0093	66.9	37 589	15 475
Y _m 495_770	97.52	−26.25	124.87	127.6	0.206	−0.0337	101.8	32 563	14 470
G _o 495_565	77.31	−135.83	89.22	162.51	0.1448	−0.0409	146.6	25 525	−1 525c
C _m 380_565	80.92	−63.4	−33.84	71.87	0.1848	−0.1061	208.0	16 480	−1 480c
B _m 380_495	30.14	96.68	−120.47	154.47	0.3221	−0.2191	308.7	11 459	28 542
M _o 575_475	74.8	82.43	−41.17	92.14	0.2629	−0.1114	333.4	−1 519c	23 519
N _o 380_770	0.9	−0.01	0.0	0.01	0.2175	−0.0887	157.8	23 515	−1 515c
W _o 380_770	100.0	0.0	0.0	0.0	0.2175	−0.0887	0.0	−1 519c	−1 515c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for D65, $Y_{w,10}=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	68.13	64.09	144.35	157.94	0.2533	0.0	66.0	37 588	13 469
Y _m 495_770	95.61	-14.82	123.91	124.79	0.2086	-0.0302	96.8	32 560	12 464
G _o 495_565	76.63	-112.27	91.34	144.73	0.1546	-0.0364	140.8	25 525	-1 525c
C _m 380_565	82.83	-64.38	-29.45	70.8	0.1826	-0.1006	204.5	15 475	-1 475c
B _m 380_495	39.48	47.92	-101.22	111.99	0.2583	-0.1769	295.3	11 459	29 546
M _o 575_475	75.51	71.82	-39.39	81.91	0.2544	-0.1072	331.2	-1 516c	23 516
N _o 380_770	0.9	-0.01	0.0	0.01	0.2152	-0.0857	158.2	22 510	-1 510c
W _o 380_770	100.0	0.0	0.0	0.0	0.2152	-0.0857	0.0	-1 516c	-1 510c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for D50, $Y_{w,10}=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	70.7	65.98	135.67	150.86	0.2547	0.0	64.0	37 589	13 469
Y _m 495_770	96.45	-8.8	110.45	110.8	0.2127	-0.0291	94.5	32 563	12 464
G _o 495_565	75.67	-111.25	77.93	135.83	0.156	-0.0357	144.9	25 525	-1 525c
C _m 380_565	80.88	-75.4	-29.92	81.12	0.1777	-0.0936	201.6	15 477	-1 477c
B _m 380_495	35.8	35.04	-97.59	103.69	0.2504	-0.1723	289.7	12 461	30 552
M _o 575_475	76.47	69.02	-33.93	76.91	0.2539	-0.0965	333.8	-1 517c	23 517
N _o 380_770	0.9	-0.01	0.0	0.01	0.2166	-0.0782	160.1	22 511	-1 511c
W _o 380_770	100.0	0.0	0.0	0.0	0.2166	-0.0782	0.0	-1 517c	-1 511c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for P40, $Y_{w,10}=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	73.9	63.66	130.14	144.87	0.2557	0.0	63.9	38 590	13 469
Y _m 495_770	97.18	-5.73	100.97	101.14	0.2178	-0.0277	93.2	33 566	12 464
G _o 495_565	73.76	-112.78	67.07	131.22	0.1575	-0.035	149.2	25 525	-1 525c
C _m 380_565	78.14	-83.58	-31.64	89.37	0.176	-0.0891	200.7	15 478	-1 478c
B _m 380_495	32.11	27.64	-95.31	99.24	0.249	-0.1713	286.1	12 461	31 557
M _o 575_475	78.27	65.12	-28.46	71.07	0.2548	-0.0874	336.3	-1 519c	23 519
N _o 380_770	0.9	-0.01	0.0	0.01	0.2203	-0.0723	161.8	22 513	-1 513c
W _o 380_770	100.0	0.0	0.0	0.0	0.2203	-0.0723	0.0	-1 519c	-1 513c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for A00, $Y_{w,10}=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	78.81	59.69	112.18	127.08	0.2584	0.0	61.9	38 592	14 470
Y _m 495_770	98.26	-0.82	77.69	77.7	0.2266	-0.0251	90.6	34 572	12 464
G _o 495_565	70.16	-114.4	44.44	122.73	0.1606	-0.0333	158.7	25 525	-1 525c
C _m 380_565	73.16	-99.94	-31.76	104.86	0.1709	-0.0769	197.6	16 481	-1 481c
B _m 380_495	25.05	6.16	-85.26	85.48	0.2344	-0.1629	274.1	12 463	34 570
M _o 575_475	81.3	58.06	-18.67	60.99	0.2567	-0.0687	342.1	-1 522c	24 522
N _o 380_770	0.9	-0.01	0.0	0.01	0.2269	-0.0591	165.3	23 518	-1 518c
W _o 380_770	100.0	0.0	0.0	0.0	0.2269	-0.0591	0.0	-1 522c	-1 518c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for E00, $Y_{w,10}=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	70.77	61.84	145.42	158.02	0.2547	0.0	66.9	37 589	13 468
Y _m 495_770	96.1	-13.37	122.07	122.8	0.2131	-0.0293	96.2	32 563	12 464
G _o 495_565	75.07	-115.82	86.82	144.75	0.1555	-0.0361	143.1	25 525	-1 525c
C _m 380_565	80.82	-68.7	-32.14	75.84	0.1836	-0.1003	205.0	15 475	-1 475c
B _m 380_495	37.42	47.17	-102.35	112.7	0.2631	-0.1794	294.7	11 459	29 549
M _o 575_475	77.05	69.39	-35.92	78.14	0.2563	-0.103	332.6	-1 517c	23 517
N _o 380_770	0.9	-0.01	0.0	0.01	0.219	-0.0837	159.0	22 510	-1 510c
W _o 380_770	100.0	0.0	0.0	0.0	0.219	-0.0837	0.0	-1 517c	-1 510c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for C00, $Y_{w,10}=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	68.69	61.05	149.19	161.2	0.2531	0.0	67.7	37 588	13 468
Y _m 495_770	95.27	-15.73	128.91	129.87	0.21	-0.0301	96.9	32 561	12 464
G _o 495_565	75.67	-112.44	94.37	146.8	0.1558	-0.0365	139.9	25 526	1 408
C _m 380_565	82.42	-61.55	-30.96	68.9	0.1858	-0.1037	206.7	14 474	-1 474c
B _m 380_495	40.81	48.35	-101.85	112.74	0.2595	-0.1776	295.3	11 459	29 546
M _o 575_475	76.47	69.51	-39.01	79.7	0.2546	-0.1091	330.6	-1 516c	23 516
N _o 380_770	0.9	-0.01	0.0	0.01	0.2171	-0.088	157.9	21 509	-1 509c
W _o 380_770	100.0	0.0	0.0	0.0	0.2171	-0.088	0.0	-1 516c	-1 509c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for P00, $Y_{w,10}=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	72.95	62.27	139.09	152.4	0.2558	0.0	65.8	38 590	13 469
Y _m 495_770	96.7	−9.13	112.34	112.71	0.2167	−0.0283	94.6	33 565	12 464
G _o 495_565	73.95	−115.37	76.76	138.58	0.1567	−0.0355	146.3	25 525	−1 525c
C _m 380_565	79.0	−76.95	−32.84	83.67	0.1803	−0.0954	203.1	15 476	−1 476c
B _m 380_495	34.6	37.77	−99.62	106.54	0.258	−0.1765	290.7	11 459	30 553
M _o 575_475	78.09	66.57	−31.57	73.67	0.2561	−0.0949	334.6	−1 518c	23 518
N _o 380_770	0.9	−0.01	0.0	0.01	0.2208	−0.0781	160.5	22 512	−1 512c
W _o 380_770	100.0	0.0	0.0	0.0	0.2208	−0.0781	0.0	−1 518c	−1 512c

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for Q00, $Y_{w,10}=100$									
<i>CodeD65</i>	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	68.52	61.0	149.86	161.8	0.2534	0.0	67.8	37 588	13 468
Y _m 495_770	95.5	−17.97	130.13	131.37	0.2093	−0.0302	97.8	32 560	12 464
G _o 495_565	76.14	−116.03	95.81	150.48	0.1544	−0.0366	140.4	25 525	−1 525c
C _m 380_565	82.55	−61.0	−30.94	68.4	0.1864	−0.1043	206.8	14 474	−1 474c
B _m 380_495	39.93	55.32	−104.07	117.86	0.2667	−0.1815	297.9	11 458	28 544
M _o 575_475	76.0	72.15	−40.09	82.54	0.2565	−0.1103	330.9	−1 516c	23 516
N _o 380_770	0.9	−0.01	0.0	0.01	0.2173	−0.0886	157.8	21 509	−1 509c
W _o 380_770	100.0	0.0	0.0	0.0	0.2173	−0.0886	0.0	−1 516c	−1 509c