

Optimal colours (o) <i>RYGCBM</i> of maximum (m) C_{AB} for D65, $Y_w=100$									
<i>Code</i> D65	L^*_{100}	a^*_{100}	b^*_{100}	C^*_{ab}	a'	b'	h_{ab}	i_d, λ^*_d	i_c, λ^*_c
R _m 565_770	70.18	61.64	132.51	146.14	0.2511	−0.0093	65.0	37 589	15 475
Y _m 495_770	97.55	−23.15	119.09	121.32	0.2052	−0.0337	101.0	32 563	14 470
G _o 495_565	77.76	−132.05	85.19	157.15	0.145	−0.0407	147.1	25 525	−1 525c
C _m 380_565	81.29	−66.98	−32.23	74.33	0.181	−0.1027	205.6	16 481	−1 481c
B _m 380_495	29.91	89.06	−117.04	147.07	0.3122	−0.2135	307.2	12 460	28 543
M _o 575_475	74.32	82.25	−40.48	91.68	0.2609	−0.1085	333.7	−1 519c	23 519
N _o 380_770	0.9	−0.01	0.0	0.01	0.2154	−0.0861	158.2	23 515	−1 515c
W _o 380_770	100.0	0.0	0.0	0.0	0.2154	−0.0861	0.0	−1 519c	−1 515c