

$XYZ_w=95.0443, 100.0, 108.89$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D65$



CIE LAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770

G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w=100$

6 de la maximum (m) C_{AB} pour D65

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w=96.4228, 100.0, 82.49$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D50$

CIELAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770


G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w=100$

6 de la maximum (m) C_{AB} pour D50

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 100.932, 100.0, 64.68$

b^*_{D65}

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P40$



CIE LAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770

G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour P40

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w = 109.849, 100.0, 35.58$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.0816$

$n = A00$

CIELAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770


G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour A00

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 100.001, 100.0, 100.0$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = E00$



CIE LAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770

G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour E00

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w=98.0718, 100.0, 118.22$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = C00$



CIE LAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770

G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w=100$

6 de la maximum (m) C_{AB} pour C00

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w = 102.067, 100.0, 81.06$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P00$



CIE LAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770

G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour P00

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w=97.9332, 100.0, 118.95$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = Q00$



$XYZ_w=94.8136, 100.0, 107.33$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D65$



CIE LAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770

G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w=100$

6 de la maximum (m) C_{AB} pour D65

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w = 96.7256, 100.0, 81.41$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D50$

CIELAB D65

Nom et la gamme spectrale

R_m 561_770 Y_m 495_770


G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour D50

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 101.751, 100.0, 64.44$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P40$

CIELAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770


G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour P40

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 111.15, 100.0, 35.19$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = A00$

CIELAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770


G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour A00

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 99.9908, 99.9999, 100.0$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = E00$



$XYZ_w = 97.2866, 100.0, 116.14$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = C00$



CIE LAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770

G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour C00

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w = 102.375, 100.0, 81.25$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P00$

CIELAB D65

Nom et la gamme spectrale

R_m 561_770 Y_m 495_770


G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour P00

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 97.65, 100.0, 118.42$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = Q00$



CIE LAB D65

Nom et la gamma spectrale

R_m 561_770 Y_m 495_770

G_m 475_573 C_m 380_561

B_m 380_495 M_m 573_475

Ostwald couleurs optimales (o), $Y_w = 100$

6 de la maximum (m) C_{AB} pour Q00

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})