

$XYZ_w=95.0443, 100.0, 108.89$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D65$

CIELAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour $D65$

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=96.4228, 100.0, 82.49$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D50$

CIELAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour $D50$

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=100.932, 100.0, 64.68$

b^*_{D65}

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.21945$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P40$



CIE LAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770

G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour P40

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w = 109.849, 100.0, 35.58$

b^*_{D65}

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = A00$



CIE LAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770

G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w = 100$

4 de la chroma maximum (m) C_{AB} pour A000

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w=100.001, 100.0, 100.0$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = E00$

CIELAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour $E00$

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=98.0718, 100.0, 118.22$ b^*

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = C00$



CIE LAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770

G_0 495_565 C_m 380_565

B_m 380_495 M_0 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour C000

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w = 102.067, 100.0, 81.06$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P00$

CIELAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w = 100$

4 de la chroma maximum (m) C_{AB} pour P0120

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=97.9332, 100.0, 118.95$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = Q00$



CIE LAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770

G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour Q000

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w=95.0443, 100.0, 108.89$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D65$

CIELAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour D65

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=96.4228, 100.0, 82.49$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D50$

CIELAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour $D50$

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=100.932, 100.0, 64.68$

b^*_{D65}

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.219451$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P40$



CIE LAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770

G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour P40

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

b* D65

$$a^* = 500 (a' - a'_n) Y^{1/3}$$

$$b^* = 500 (b' - b'_n) Y^{1/3}$$


$$a' = a_2 [x/y]^{1/3}$$

$$b' = b_2 [z/y]^{1/3}$$

$$a_2 = [1/X_{D65}]^{1/3} = 0.2191$$

$$b_7 = -[1/Z_{D65}]^{1/3} = -0.98376$$

$$M = A00$$



CIELAB D65

Nom et la gamme spectrale

$$R_m \ 565_770 \quad Y_m \ 495_770$$

G_0 495_565 C_m 380_565

 B_m 380_495 M_0 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour A020

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w=100.001, 100.0, 100.0$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = E00$

CIELAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour $E00$

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=98.0718, 100.0, 118.22$ b^*

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = C00$



CIE LAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770

G_0 495_565 C_m 380_565

B_m 380_495 M_0 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour $C00$

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})

$XYZ_w = 102.067, 100.0, 81.06$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P00$

CIELAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w = 100$

4 de la chroma maximum (m) C_{AB} pour P0120

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=97.9332, 100.0, 118.95$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = Q00$



CIE LAB D65

Nom et la gamma spectrale

R_m 565_770 Y_m 495_770

G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Couleurs optimales (o), $Y_w=100$

4 de la chroma maximum (m) C_{AB} pour $Q000$

dans le diagramme de chroma (a^*_{D65}, b^*_{D65})