

$XYZ_w = 84.1998, 88.59, 96.46$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2191$

$b_2 = -[1/Z_n]^{1/3} = -0.08376$

$n = D65$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per D65

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w=85.421, 88.59, 73.08$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.218$

$b_2 = -[1/Z_n]^{1/3} = -0.09188$

$n = D50$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w=88,6$

4 di massimo (m) C_{AB} per D50

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 89.4154, 88.59, 57.3$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2147$

$b_2 = -[1/Z_n]^{1/3} = -0.09964$

$n = P40$

CIELAB 76

Nome e la gamma spettrali

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per P40

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w=97.3152, 88.59, 31.52$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2088$

$b_2 = -[1/Z_n]^{1/3} = -0.12161$

$n = A00$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w=88,6$

4 di massimo (m) C_{AB} per A00

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 88.5907, 88.59, 88.59$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2154$

$b_2 = -[1/Z_n]^{1/3} = -0.08617$

$n = E00$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per E00

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 86.8818, 88.59, 104.73$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2168$

$b_2 = -[1/Z_n]^{1/3} = -0.08149$

$n = C00$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per C00

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w=90.421, 88.59, 71.81$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2139$

$b_2 = -[1/Z_n]^{1/3} = -0.09242$

$n = P00$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w=88,6$

4 di massimo (m) C_{AB} per P00

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w=86.7591, 88.59, 105.38$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2169$

$b_2 = -[1/Z_n]^{1/3} = -0.08133$

$n = Q00$



CIE LAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770

G_0 495_565 C_m 380_565

B_m 380_495 M_0 565_495

Colori ottimi (o), $Y_w=88,6$

4 di massimo (m) C_{AB} per $Q00$

nel diagramma di CIE LAB (a^* , b^*)

$XYZ_w = 84.1998, 88.59, 96.46$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2191$

$b_2 = -[1/Z_n]^{1/3} = -0.08376$

$n = D65$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per D65

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w=85.421, 88.59, 73.08$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.218$

$b_2 = -[1/Z_n]^{1/3} = -0.09188$

$n = D50$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w=88,6$

4 di massimo (m) C_{AB} per D50

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 89.4154, 88.59, 57.3$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2147$

$b_2 = -[1/Z_n]^{1/3} = -0.09964$

$n = P40$

CIELAB 76

Nome e la gamma spettrali

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per P40

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w=97.3152, 88.59, 31.52$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2088$

$b_2 = -[1/Z_n]^{1/3} = -0.12161$

$n = A00$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w=88,6$

4 di massimo (m) C_{AB} per A00

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 88.5907, 88.59, 88.59$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2154$

$b_2 = -[1/Z_n]^{1/3} = -0.08617$

$n = E00$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per E00

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 86.8818, 88.59, 104.73$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2168$

$b_2 = -[1/Z_n]^{1/3} = -0.08149$

$n = C00$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per C00

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w=90.421, 88.59, 71.81$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2139$

$b_2 = -[1/Z_n]^{1/3} = -0.09242$

$n = P00$

CIELAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w=88,6$

4 di massimo (m) C_{AB} per P00

nel diagramma di CIELAB (a^*, b^*)



$XYZ_w = 86.7591, 88.59, 105.38$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2169$

$b_2 = -[1/Z_n]^{1/3} = -0.08133$

$n = Q00$



CIE LAB 76

Nome e la gamma spettrale

R_m 565_770 Y_m 495_770

G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Colori ottimi (o), $Y_w = 88,6$

4 di massimo (m) C_{AB} per $Q00$

nel diagramma di CIE LAB (a^*, b^*)