



### Mathematical equations of hyperbel functions

See: Papula, L., (2003), Mathematische Formelsammlung, Vieweg

$$F(x) = \tanh(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}} = \frac{u(x)}{v(x)} \quad u'(x) = v(x) \quad [1]$$

$$F'(x) = \frac{u'(x)v(x) - u(x)v'(x)}{v^2(x)} = \frac{v^2(x) - u^2(x)}{v^2(x)} \quad [2]$$

$$F'(x) = \frac{[e^x + e^{-x}][e^x - e^{-x}] - [e^x - e^{-x}][e^x - e^{-x}]}{[e^x + e^{-x}]^2} \quad [3]$$

$$F'(x) = \frac{4}{[e^x + e^{-x}]^2} = \frac{1}{\cosh^2(x)} \quad [4]$$

eeo40-1n eeo31-2n

### Mathematical equations of hyperbel functions

See: Papula, L., (2003), Mathematische Formelsammlung, Vieweg

$$F(x/a) = \tanh(x/a) = \frac{e^{x/a} - e^{-x/a}}{e^{x/a} + e^{-x/a}} = \frac{u(x/a)}{v(x/a)} \quad [1]$$

$$F'(x/a) = \frac{u'(x/a)v(x/a) - u(x/a)v'(x/a)}{v^2(x/a)} \quad [2]$$

$$F'(x/a) = \frac{v^2(x/a) - u^2(x/a)}{av^2(x/a)} \quad [3]$$

$$F'(x/a) = \frac{4}{a[e^{x/a} + e^{-x/a}]^2} = \frac{1}{a \cosh^2(x/a)} \quad [4]$$

eeo40-2n eeo31-4n

$F_{ab}(x_r) = \text{achromatic receptor response}$

$$F_{ab}(x_r) = b \frac{e^{x_r/a} - e^{-x_r/a}}{e^{x_r/a} + e^{-x_r/a}} \quad a=1.00, b=1.00, c=2.718282$$

$$F'_{ab}(x_r) = b \frac{4b}{(e^{x_r/a} + e^{-x_r/a})^2} = b \frac{4b}{(e^{x_r/a} + e^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$\text{Asymptote } F'_{ab}(x_r) = 4b \cdot \frac{e^{x_r/a} - e^{-x_r/a}}{(e^{x_r/a} + e^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$m_{1,01}=0.98 \quad N \quad W \quad F_{1,-1}(x_r)$$

$$L_u = 28 \text{ cd/m}^2 \quad L_u = \log(L/L_u) \quad \text{range of office illuminances}$$



### Mathematical equations of hyperbel functions

See: Papula, L., (2003), Mathematische Formelsammlung, Vieweg

$$F_{ab}(x/a) = b \tanh(x/a) = b \frac{e^{x/a} - e^{-x/a}}{e^{x/a} + e^{-x/a}} = b \frac{u(x/a)}{v(x/a)} \quad [1]$$

$$F'_{ab}(x/a) = b \frac{u'(x/a)v(x/a) - u(x/a)v'(x/a)}{v^2(x/a)} \quad [2]$$

$$F'_{ab}(x/a) = b \frac{v^2(x/a) - u^2(x/a)}{av^2(x/a)} \quad [3]$$

$$F'_{ab}(x/a) = \frac{4b}{a[e^{x/a} + e^{-x/a}]^2} = \frac{b}{a \cosh^2(x/a)} \quad [4]$$

eeo41-2n

$F_{ab}(x_r) = \text{achromatic receptor response}$

$$F_{ab}(x_r) = b \frac{e^{x_r/a} - e^{-x_r/a}}{e^{x_r/a} + e^{-x_r/a}} \quad a=1.00, b=1.00, c=2.718282$$

$$F'_{ab}(x_r) = b \frac{4b}{(e^{x_r/a} + e^{-x_r/a})^2} = b \frac{4b}{(e^{x_r/a} + e^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$\text{Asymptote } F'_{ab}(x_r) = 4b \cdot \frac{e^{x_r/a} - e^{-x_r/a}}{(e^{x_r/a} + e^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$m_{1,01}=0.98 \quad N \quad W \quad F_{1,-1}(x_r)$$

$$L_u = 28 \text{ cd/m}^2 \quad L_u = \log(L/L_u) \quad \text{range of office illuminances}$$



### Mathematical equations of hyperbel functions

See: Papula, L., (2003), Mathematische Formelsammlung, Vieweg

$$F_{ab}(x/a) = b \tanh(x/a) = b \frac{e^{x/a} - e^{-x/a}}{e^{x/a} + e^{-x/a}} = b \frac{u(x/a)}{v(x/a)} \quad [1]$$

$$F'_{ab}(x/a) = b \frac{u'(x/a)v(x/a) - u(x/a)v'(x/a)}{v^2(x/a)} \quad [2]$$

$$F'_{ab}(x/a) = b \frac{v^2(x/a) - u^2(x/a)}{av^2(x/a)} \quad [3]$$

$$F'_{ab}(x/a) = \frac{4b}{a[e^{x/a} + e^{-x/a}]^2} = \frac{b}{a \cosh^2(x/a)} \quad [4]$$

eeo41-4n

$F_{ab}(x_r) = \text{achromatic receptor response}$

$$F_{ab}(x_r) = b \frac{e^{x_r/a} - e^{-x_r/a}}{e^{x_r/a} + e^{-x_r/a}} \quad a=1.00, b=1.00, c=2.718282$$

$$F'_{ab}(x_r) = b \frac{4b}{(e^{x_r/a} + e^{-x_r/a})^2} = b \frac{4b}{(e^{x_r/a} + e^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$\text{Asymptote } F'_{ab}(x_r) = 4b \cdot \frac{e^{x_r/a} - e^{-x_r/a}}{(e^{x_r/a} + e^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$m_{1,01}=0.98 \quad N \quad W \quad F_{1,-1}(x_r)$$

$$L_u = 28 \text{ cd/m}^2 \quad L_u = \log(L/L_u) \quad \text{range of office illuminances}$$



$F_{ab}(x_r) = \text{achromatic receptor response}$

$$F_{ab}(x_r) = b \frac{10^{x_r/a} - 10^{-x_r/a}}{10^{x_r/a} + 10^{-x_r/a}} = b \frac{10^{x_r/a} - 10^{-x_r/a}}{10^{x_r/a} + 10^{-x_r/a}} \quad a=1.00, b=1.00, c=2.718282$$

$$F'_{ab}(x_r) = b \frac{4b}{(10^{x_r/a} + 10^{-x_r/a})^2} = b \frac{4b}{(10^{x_r/a} + 10^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$\text{Asymptote } F'_{ab}(x_r) = 4b \cdot \frac{10^{x_r/a} - 10^{-x_r/a}}{(10^{x_r/a} + 10^{-x_r/a})^2} = 4b \cdot \frac{10^{x_r/a} - 10^{-x_r/a}}{(10^{x_r/a} + 10^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$m_{1,01}=0.98 \quad N \quad W \quad F_{1,-1}(x_r)$$

$$L_u = 28 \text{ cd/m}^2 \quad L_u = \log(L/L_u) \quad \text{range of office illuminances}$$



$F_{ab}(x_r) = \text{achromatic receptor response}$

$$F_{ab}(x_r) = b \frac{10^{x_r/a} - 10^{-x_r/a}}{10^{x_r/a} + 10^{-x_r/a}} = b \frac{10^{x_r/a} - 10^{-x_r/a}}{10^{x_r/a} + 10^{-x_r/a}} \quad a=1.00, b=1.00, c=2.718282$$

$$F'_{ab}(x_r) = b \frac{4b}{(10^{x_r/a} + 10^{-x_r/a})^2} = b \frac{4b}{(10^{x_r/a} + 10^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$\text{Asymptote } F'_{ab}(x_r) = 4b \cdot \frac{10^{x_r/a} - 10^{-x_r/a}}{(10^{x_r/a} + 10^{-x_r/a})^2} = 4b \cdot \frac{10^{x_r/a} - 10^{-x_r/a}}{(10^{x_r/a} + 10^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$m_{1,01}=0.98 \quad N \quad W \quad F_{1,-1}(x_r)$$

$$L_u = 28 \text{ cd/m}^2 \quad L_u = \log(L/L_u) \quad \text{range of office illuminances}$$



$F_{ab}(x_r) = \text{achromatic receptor response}$

$$F_{ab}(x_r) = b \frac{10^{x_r/a} - 10^{-x_r/a}}{10^{x_r/a} + 10^{-x_r/a}} = b \frac{10^{x_r/a} - 10^{-x_r/a}}{10^{x_r/a} + 10^{-x_r/a}} \quad a=1.00, b=1.00, c=2.718282$$

$$F'_{ab}(x_r) = b \frac{4b}{(10^{x_r/a} + 10^{-x_r/a})^2} = b \frac{4b}{(10^{x_r/a} + 10^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$\text{Asymptote } F'_{ab}(x_r) = 4b \cdot \frac{10^{x_r/a} - 10^{-x_r/a}}{(10^{x_r/a} + 10^{-x_r/a})^2} = 4b \cdot \frac{10^{x_r/a} - 10^{-x_r/a}}{(10^{x_r/a} + 10^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$m_{1,01}=0.98 \quad N \quad W \quad F_{1,-1}(x_r)$$

$$L_u = 28 \text{ cd/m}^2 \quad L_u = \log(L/L_u) \quad \text{range of office illuminances}$$



$F_{ab}(x_r) = \text{achromatic receptor response}$

$$F_{ab}(x_r) = b \frac{10^{x_r/a} - 10^{-x_r/a}}{10^{x_r/a} + 10^{-x_r/a}} = b \frac{10^{x_r/a} - 10^{-x_r/a}}{10^{x_r/a} + 10^{-x_r/a}} \quad a=1.00, b=1.00, c=2.718282$$

$$F'_{ab}(x_r) = b \frac{4b}{(10^{x_r/a} + 10^{-x_r/a})^2} = b \frac{4b}{(10^{x_r/a} + 10^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$\text{Asymptote } F'_{ab}(x_r) = 4b \cdot \frac{10^{x_r/a} - 10^{-x_r/a}}{(10^{x_r/a} + 10^{-x_r/a})^2} = 4b \cdot \frac{10^{x_r/a} - 10^{-x_r/a}}{(10^{x_r/a} + 10^{-x_r/a})^2} \quad a=1.00, b=1.00$$

$$m_{1,01}=0.98 \quad N \quad W \quad F_{1,-1}(x_r)$$

$$L_u = 28 \text{ cd/m}^2 \quad L_u = \log(L/L_u) \quad \text{range of office illuminances}$$



see similar files of the whole serie: <http://farbe.li.tu-berlin.de> or <http://color.li.tu-berlin.de>

technical information: <http://farbe.li.tu-berlin.de/eeosh.htm>

TUB-test chart eeo4; Model of normalized response function  $F_{ab}(x_r)$  and derivation  $F'_{ab}(x_r)$

Mathematical calculation of the derivation  $F'_{ab}(x_r)$ , of the contrast  $L/\Delta L$ , and the discrimination  $\Delta L$

see similar files: <http://farbe.li.tu-berlin.de/eeo4l0na.txt /ps>