

Achromatisches Sehen mit relativer Leuchtdichte

Mathematische Gleichungen mit Potenzfunktionen

$$F_{cb}(L_R, n) = b \tanh(x_r/c) = b \frac{L_R^n - L_R^{-n}}{L_R^n + L_R^{-n}} \quad \begin{array}{l} x_r = \log(L_R) \\ L_R = L/L_u \\ x_r \geq 0 \end{array} \quad [1]$$

$$\frac{dF_{cb}(L_R, n)}{dL_R} = \frac{4bn}{L_R[L_R^n + L_R^{-n}]^2} \quad \begin{array}{l} x_r = \ln L_R / \ln(10) \\ dx_r/dL_R = 1/(\ln(10)L_R) \\ n = 1/(\ln(10)c) \end{array} \quad [5]$$

$$\frac{L}{dL} = \frac{4bnL_u L}{L_R[L_R^n + L_R^{-n}]^2} \quad dL = \frac{L_R[L_R^n + L_R^{-n}]^2}{4bnL_u} \quad [7]$$

$$\frac{L/dL}{(L/dL)_u} = \frac{4L}{L_R[L_R^n + L_R^{-n}]^2 L_u}; \quad \frac{dL}{dL_u} = \frac{L_R[L_R^n + L_R^{-n}]^2}{4} \quad [8]$$